

ON THE DIAGRAM OF 132-AVOIDING PERMUTATIONS

Astrid Reifegerste

Institut für Mathematik, Universität Hannover

Welfengarten 1

D-30167 Hannover, Germany

reifegerste@math.uni-hannover.de

October 15, 2002

ABSTRACT. The diagram of a 132-avoiding permutation can easily be characterized: it is simply the diagram of a partition. Based on this fact, we present a new bijection between 132-avoiding and 321-avoiding permutations. We will show that this bijection translates the correspondences between these permutations and Dyck paths given by Krattenthaler and by Billey-Jockusch-Stanley, respectively, to each other. Moreover, the diagram approach yields simple proofs for some enumerative results concerning forbidden patterns in 132-avoiding permutations.

1 INTRODUCTION

Let $\mathcal{S}_n$ denote the symmetric group on $\{1, \dots, n\}$. Given a permutation $\pi = \pi_1 \cdots \pi_n \in \mathcal{S}_n$ and a permutation $\tau = \tau_1 \cdots \tau_k \in \mathcal{S}_k$, we say that π *contains the pattern* τ if there is a sequence $1 \leq i_1 < i_2 < \dots < i_k \leq n$ such that the elements $\pi_{i_1} \pi_{i_2} \cdots \pi_{i_k}$ are in the same relative order as $\tau_1 \tau_2 \cdots \tau_k$. Otherwise, π *avoids the pattern* τ , or alternatively, π is τ -*avoiding*. We denote by $\mathcal{S}_n(\tau)$ the set of all permutations in $\mathcal{S}_n$ which avoid τ .

It is an often quoted fact that $|\mathcal{S}_n(\tau)|$ is equal to the n th Catalan number $C_n = \frac{1}{n+1} \binom{2n}{n}$ for each pattern $\tau \in \mathcal{S}_3$. Because of obvious symmetry arguments, from an enumerative viewpoint there are only two distinct cases to consider, $\tau \in \{123, 321\}$ and $\tau \in \{132, 213, 231, 312\}$. Several authors established bijections between permutations avoiding a pattern of each of these classes. The first one was given by Simion and Schmidt [17]; West described in [18] a construction using trees; and recently, Krattenthaler [9] connected the 123-avoiding and 132-avoiding permutations via Dyck paths.

In Section 2, we present a bijection between $\mathcal{S}_n(321)$ and $\mathcal{S}_n(132)$ based on another interesting combinatorial object, the diagrams.

Our construction has the advantage that the excedances of a permutation in $\mathcal{S}_n(321)$ are precisely the descents of its image in $\mathcal{S}_n(132)$.

An *excedance* of π is an integer $i \in \{1, \dots, n-1\}$ such that $\pi_i > i$. Here the element π_i is called an *excedance letter* for π . Given a permutation π , we denote the set of excedances of π by $E(\pi)$ and the number $|E(\pi)|$ by $\text{exc}(\pi)$. An integer $i \in \{1, \dots, n-1\}$ for which $\pi_i > \pi_{i+1}$ is called a *descent* of π . If i is a descent, we say that π_{i+1} is a *descent bottom* for π . The set of descents of π is denoted by $D(\pi)$, its cardinality is denoted by $\text{des}(\pi)$, as usual.

Given a permutation $\pi \in \mathcal{S}_n$, let $i_1 < i_2 < \dots < i_e$ be the excedances of π and $j_1 < j_2 < \dots < j_{n-e}$ the remaining positions. It characterizes 321-avoiding permutations that both subwords $\pi_{i_1}\pi_{i_2}\dots\pi_{i_e}$ and $\pi_{j_1}\pi_{j_2}\dots\pi_{j_{n-e}}$ are increasing. Thus any 321-avoiding permutation is uniquely determined by its excedances and excedance letters.

There are several correspondences between restricted permutations and lattice paths, in particular, Dyck paths. A *Dyck path* is a path in the (x, y) -plane from the origin to $(2n, 0)$ with steps $[1, 1]$ (called up-steps) and $[1, -1]$ (called down-steps) that never falls below the x -axis.

For 321-avoiding permutations such a bijection was given by Billey, Jockusch and Stanley [1]; for 132-avoiding permutations Krattenthaler proposed a correspondence to Dyck paths in [9]. In Section 3 we will show that the Dyck path obtained for any $\pi \in \mathcal{S}_n(321)$ by the first mentioned correspondence and the Dyck path associating by Krattenthaler's correspondence with the image (with respect to our bijection) $\sigma \in \mathcal{S}_n(132)$ of π are the same.

Moreover, it will turn out that the diagram of a 132-avoiding permutation is closely related to the corresponding Dyck path. This yields a simple explanation for the connections between the number of inversions of the permutation and several parameters of the Dyck path recently appeared in [2]. Further, we use this relation to enumerate restricted partitions of prescribed rank.

In Section 4 the diagram approach will be used to obtain some enumerative results concerning the restriction of 132-avoiding permutations by additional patterns. These results are already known (see [11]) but we will give a bijective proof of them.

The paper ends with a note on how to obtain the number of occurrences of the pattern 132 in an arbitrary permutation via the diagram.


2 A BIJECTION BETWEEN 132-AVOIDING AND 321-AVOIDING PERMUTATIONS

Let $\mathcal{Y}_n := \{(\lambda_1, \dots, \lambda_{n-1}) : 0 \leq \lambda_{n-1} \leq \lambda_{n-2} \leq \dots \leq \lambda_1 \leq n-1, \lambda_i \leq n-i \text{ for all } i\}$ be the set of partitions whose Young diagram fits in the shape $(n-1, n-2, \dots, 1)$. (We will identify a partition with its Young diagram and vice versa.) In [15, Sect. 3.2] (or [16]), we describe a bijection $\mathcal{S}_n(321) \rightarrow \mathcal{Y}_n$ which takes the permutation π with excedances $i_1, \dots, i_e$ to the diagram with corners $(i_k, n+1-\pi_{i_k})$ where $k = 1, \dots, e$. For 132-avoiding permutations a

simple correspondence to partitions with restricted diagram can be given, as well.

The key object in our derivation is the *diagram of a permutation* (for an introduction see [10, chap. 1]). Given a permutation $\pi \in \mathcal{S}_n$, we obtain the diagram $D(\pi)$ of π as follows. Let π be represented by an $n \times n$ -array with a dot in each of the squares (i, π_i) . (The other cells are white.) Shadow all squares due south or due east of some dot and the dotted cell itself. The diagram $D(\pi)$ is defined as the region left unshaded after this procedure.

Example 2.1 The diagram of $\pi = 4\ 2\ 8\ 3\ 6\ 9\ 7\ 5\ 1\ 10 \in \mathcal{S}_{10}$ contains the white squares of




By the construction, each of the connected components of $D(\pi)$ is a Young diagram. Their corners are defined to be the elements of the *essential set* $\mathcal{E}(\pi)$ of the permutation π . In [8], Fulton introduced this set which together with a rank function was used as a tool for algebraic treatment of Schubert polynomials. In [5], Eriksson and Linusson characterized the essential sets that can arise from arbitrary permutations, as well as those coming from certain classes of permutations.

It is very easy to characterize the diagrams of 132-avoiding permutations.

Theorem 2.2 *Let $\pi \in \mathcal{S}_n$ be a permutation not equal to the identity. Then π is 132-avoiding if and only if its diagram consists of only one component and $(1, 1) \in D(\pi)$.*

Proof. If there are indices $i < j < k$ such that $\pi_i < \pi_k < \pi_j$, then the square (j, π_k) belongs to $D(\pi)$, but it is not connected with $(1, 1)$:



Clearly, the existence of such a square is also sufficient for π containing the pattern 132.

Note that the square $(1, 1)$ must be an element of $D(\pi)$ for any 132-avoiding permutation $\pi \neq \text{id}$, otherwise we would have $\pi_1 = 1$ and hence $\pi_i = i$ for all $i = 1, \dots, n$. $\square$


Thus the diagram $D(\pi)$ of a permutation $\pi \in \mathcal{S}_n(132)$ is the graphical representation of a partition. By construction, $D(\pi)$ is just the diagram of an element of $\mathcal{Y}_n$: the square $(i, j(i))$ belongs

to $D(\pi)$ if and only if no index $k \leq i$ satisfies $\pi_k \leq j$. Thus we have $j(i) \leq n - i$.

This yields a simple bijection between 321-avoiding and 132-avoiding permutations on $\{1, \dots, n\}$ which is in the following denoted by Φ .

Let $i_1, \dots, i_e$ be the excedances of $\pi \in \mathcal{S}_n(321)$. Then the diagram of the corresponding permutation $\Phi(\pi)$ is equal to the Young diagram with corners $(i_k, n+1-\pi_{i_k})$, $k = 1, \dots, e$. Recovering $\Phi(\pi)$ from $D(\Phi(\pi))$ is trivial: row by row, put a dot in the leftmost shaded square such that there is exactly one dot in each column.

Example 2.3 For the permutation $\pi = 14723856109 \in \mathcal{S}_{10}(321)$ we have $E(\pi) = \{2, 3, 6, 9\}$. Hence it corresponds to the permutation with the diagram



and we obtain $\Phi(\pi) = 89546723101 \in \mathcal{S}_{10}(132)$.

As observed by Fulton in [8], every row of a permutation diagram containing a white corner (that is an element of the essential set) corresponds to a descent. Thus we have $\text{des}(\Phi(\pi)) = \text{exc}(\pi)$ for all $\pi \in \mathcal{S}_n(321)$. But there is more to it than that: the excedance set of π and the descent set of $\Phi(\pi)$ have not only the same number of elements; the sets are even identical.

Proposition 2.4 *We have $E(\pi) = D(\Phi(\pi))$ for all $\pi \in \mathcal{S}_n(321)$.*

Proof. Any excedance i of π corresponds to a corner $(i, n+1-\pi_i)$ of $D(\Phi(\pi))$. Obviously, by constructing $\Phi(\pi)$ from its diagram we obtain a descent of $\Phi(\pi)$ at the position i . $\square$

Every 321-avoiding permutation is completely determined by its excedances and excedance letters. Our bijection shows that it is sufficient for fixing a 132-avoiding permutation to know the descents, the descent bottoms, and the first letter.

Let $i_1 < \dots < i_e$ be the excedances of $\pi \in \mathcal{S}_n(321)$, and let $\sigma := \Phi(\pi)$. Then we have

$$\begin{aligned} \sigma_1 &= n+2-\pi_{i_1}, \\ \sigma_{i_k+1} &= n+2-\pi_{i_{k+1}} \quad \text{for } k = 1, \dots, e-1, \\ \sigma_{i_e+1} &= 1. \end{aligned}$$

It is clear from the construction that these elements are precisely the left-to-right minima of σ . (A *left-to-right minimum* of a permutation σ is an element σ_i which is smaller than all elements

to its left, i.e., $\sigma_i < \sigma_j$ for every $j < i$.) Based on this, we can determine the permutation σ since σ is 132-avoiding.

Example 2.5 Let again $\pi = 1 \underline{4} \underline{7} 2 3 \underline{8} 5 6 \underline{10} 9 \in \mathcal{S}_{10}(321)$. (The underlined positions are just the excedances of π .) As described above, we obtain the left-to-right minima of $\Phi(\pi)$ and their positions:

$$8 * 5 4 * * 2 * * 1,$$

and hence, by putting the remaining elements $a = 3, 6, 7, 9, 10$ on the first possible position following $a - 1$, the permutation $\Phi(\pi) = 8 9 5 4 6 7 2 3 10 1$.

Remarks 2.6

a) In Chapter 1 of [10], Macdonald defined the *dominant* permutations. This special case of *vexillary* (or 2143-avoiding) permutations is characterized by the following equivalent conditions:

- (i) $D(\pi)$ is the diagram of a partition.
- (ii) The code c of π is a partition, that is, $c_i \geq c_{i+1}$ for all $i = 1, \dots, n - 1$.

(For a permutation π the i th component of its *code* counts the number of indices $j > i$ satisfying $\pi_j < \pi_i$.) As mentioned above, we may assume in (i) that $D(\pi)$ is an element of $\mathcal{Y}_n$. Thus $\mathcal{S}_n(132)$ is precisely the set of all dominant permutations. In particular, there are C_n such permutations.

- b) For *any* permutation $\pi \in \mathcal{S}_n$ the number of squares in the i th row of $D(\pi)$ is equal to the i th component c_i of its code (see [10, p. 9]). Hence it follows that $|D(\pi)| = c_1 + \dots + c_n = \text{inv}(\pi)$, where $\text{inv}(\pi)$ denotes the number of inversions of π .
- c) In [15, Sect. 3.2] (or in [16]), we prove that the number of excedances has the Narayana distribution over $\mathcal{S}_n(321)$. Consequently, the number of diagrams fitting in the shape $(n - 1, n - 2, \dots, 1)$ and having k corners is given by the Narayana number $N(n, k + 1) = \frac{1}{n} \binom{n}{k} \binom{n}{k+1}$. Thus the bijection immediately shows that the statistic **des** is Narayana distributed over $\mathcal{S}_n(132)$.
- d) Also in [15, Sect. 2.3], an involution on $\mathcal{S}_n(321)$ was established which proves the symmetry of the joint distribution of the pair **(exc, inv)** over $\mathcal{S}_n(321)$. For $\pi \in \mathcal{S}_n(321)$ let $\sigma = \sigma_1 \cdots \sigma_n$ be the image with respect to this map. Then the reverse $\sigma_n \cdots \sigma_1 \in \mathcal{S}_n(123)$ corresponds to $\Phi(\pi) \in \mathcal{S}_n(132)$ by the bijection due to Simion and Schmidt (see [17, Prop. 19]).
- e) In [7, p. 7], Fulmek has given a "graphical" construction of a map from $\mathcal{S}_n$ to the set of Dyck paths of length $2n$ whose restrictions on $\mathcal{S}_n(321)$ and $\mathcal{S}_n(312)$, respectively, are bijections. Based on the graph of a permutation $\pi \in \mathcal{S}_n$ he considered the union of all points which are in the southeast of some left-to-right maximum. The corresponding path

is defined to be the boundary of this region. It is easy to see that the region above the path is just the connected component of all squares of rank zero belonging to the diagram of the inverse of $\pi_n \pi_{n-1} \cdots \pi_1$. (See also [5] for a description how to obtain an arbitrary permutation from its ranked essential set.)

3 CORRESPONDENCES TO DYCK PATHS

For 321-avoiding, as well as for 132-avoiding permutations one-to-one correspondences to lattice paths were given by several authors. In [1, p. 361], Billey, Jockusch and Stanley established a bijection Ψ_{BJS} between 321-avoiding permutations on $\{1, \dots, n\}$ and Dyck paths of length $2n$. Recently in [9, Sect. 2], Krattenthaler exhibited a Dyck path correspondence Ψ_K for 132-avoiding permutations. Our bijection Φ translates these constructions into each other.

Theorem 3.1 *Let $\pi \in \mathcal{S}_n(321)$. Then we have $\Psi_{BJS}(\pi) = \Psi_K(\Phi(\pi))$.*

Proof. Let $\pi \in \mathcal{S}_n(321)$ with the excedances $i_1 < \dots < i_e$, and let $\sigma := \Phi(\pi)$. The bijection Ψ_{BJS} constructs the Dyck path corresponding to π as follows:

- 1) Let $a_k := \pi_{i_k} - 1$ for $k = 1, \dots, e$ and $a_0 := 0$, $a_{e+1} := n$. Furthermore, let $b_k := i_k$ for $k = 1, \dots, e$ and $b_0 := 0$, $b_{e+1} := n$.
- 2) Generate the Dyck path (starting at the origin) by adjoining $a_k - a_{k-1}$ up-steps and $b_k - b_{k-1}$ down-steps for $k = 1, \dots, e + 1$.

As shown in the preceding section, the elements

$$\begin{aligned} c_1 &:= \sigma_1 = n + 2 - \pi_{i_1}, \\ c_{k+1} &:= \sigma_{i_k+1} = n + 2 - \pi_{i_{k+1}} \quad \text{for } k = 1, \dots, e - 1, \\ c_{e+1} &:= \sigma_{i_e+1} = 1 \end{aligned}$$


are the left-to-right minima of σ . With the convention $c_0 := n + 1$ we have $c_{k-1} - c_k = a_k - a_{k-1}$ for all $k = 1, \dots, e + 1$. For the number d_k of the positions between the k th and (including) the $(k + 1)$ st left-to-right minimum we obtain $d_k = b_k - b_{k-1}$ for $k = 1, \dots, e + 1$. (Let $n + 1$ be the position of the imaginary $(e + 2)$ nd minimum, so $d_{e+1} = n - b_e$.)

Hence the translation of Ψ_{BJS} by Φ constructs the Dyck path corresponding to $\sigma \in \mathcal{S}_n(132)$ as follows:

- 1) Let $c_1 > \dots > c_{e+1}$ be the left-to-right minima of σ . Furthermore, let d_k be one plus the number of the letters in σ between c_k and c_{k+1} for $k = 1, \dots, e + 1$. Initialize $c_0 := n + 1$.
- 2) Generate the Dyck path (starting at the origin) by adjoining $c_{k-1} - c_k$ up-steps and d_k down-steps for $k = 1, \dots, e + 1$.

But this is precisely the description of Ψ_K proposed in [9]. $\square$


Example 3.2 Let $\pi = 1\ 4\ 7\ 2\ 3\ 8\ 5\ 6\ 10\ 9 \in \mathcal{S}_{10}(321)$, and let $\sigma = \Phi(\pi) = 8\ 9\ 5\ 4\ 6\ 7\ 2\ 3\ 10\ 1$. Billey-Jockusch-Stanley's bijection takes π to the Dyck path



which is exactly the path corresponding to σ by Krattenthaler's bijection.

The following results use the (now obvious) fact that the Dyck path $\Psi_K(\pi)$ and the diagram of a 132-avoiding permutation π are closely related to each other. Given a permutation $\pi \in \mathcal{S}_n(132)$, its diagram $D(\pi)$ is just the region bordered by the lines between the lattice points $(0,0)$ and (n,n) and between (n,n) and $(2n,0)$, respectively, and the path $\Psi_K(\pi)$. (The northwest-to-southeast diagonals correspond to the diagram columns.)

Example 3.3 For $\pi = 8\ 9\ 5\ 4\ 6\ 7\ 2\ 3\ 10\ 1 \in \mathcal{S}_{10}(132)$ we obtain:




In [2], the authors studied the statistic e_k that counts the number of increasing subsequences of length $k + 1$ in a permutation. Expressing $e_k(\pi)$ in terms of the Dyck path $\Psi_K(\pi)$, some applications were given in [2] which relate various combinatorial structures to 132-avoiding permutations. The translation of the statistic into Dyck path characteristics becomes now immediately clear.

Corollary 3.4 *For any permutation $\pi \in \mathcal{S}_n(132)$ consider the associated Dyck path $\Psi_K(\pi)$ and denote the height of the starting point of its i th step by $w_i(\pi)$. Then we have*


- a) $w_1(\pi) + \dots + w_{2n}(\pi) = n^2 - 2 \operatorname{inv}(\pi)$.
- b) $w_{i_1}(\pi) + \dots + w_{i_s}(\pi) = \binom{n+1}{2} - \operatorname{inv}(\pi)$, where $i_1, \dots, i_s$ are the indices of the down-steps.

Proof. As remarked in 2.6b), we have $|D(\pi)| = \operatorname{inv}(\pi)$ for all $\pi \in \mathcal{S}_n$.

a) The sum of heights of the Dyck path $(\nearrow^n \searrow^n)$ (where exponentiation denotes repetition) is equal to n^2 . Any square of $D(\pi)$ reduces this value by 2:



b) The sum of all down-step heights counts the number of squares of the slanting lattices below the path:



For the path $(\nearrow)^n(\searrow)^n$ this number is $\binom{n+1}{2}$. $\square$

Remarks 3.5

- a) The sum of the heights of all steps is easily seen to equal the area of the Dyck path. It was shown in [2, Sect. 3.1] that the sum relates to the statistic $2e_1 + e_0$ for 132-avoiding permutations.
- b) For a correspondence between fountains of coins and 132-avoiding permutations, [2] used that $e_1(\pi) + e_0(\pi)$ is equal to the sum of the heights of the down-steps in $\Psi_K(\pi)$.

On the other hand, from [2] we can derive enumerative results for the partitions in $\mathcal{Y}_n$ and (using the bijection Φ) for 321-avoiding permutations.

By [2, Prop. 7], the distribution of right-to-left maxima in $\mathcal{S}_n(132)$ is given by the ballot numbers. (A *right-to-left maxima* of a permutation π is an element π_i which is larger than all π_j with $j > i$.) The number of permutations in $\mathcal{S}_n(132)$ with k such maxima equals the ballot number

$$b(n-1, n-k) = \frac{k}{2n-k} \binom{2n-k}{n}.$$

Among other things, the *ballot number* $b(n, k)$ counts the number of the lattice paths from $(0, 0)$ to $(n+k, n-k)$ with up-steps and down-steps only, never falling below the x -axis. It is well known that $b(n, k) = \binom{n+k}{n} - \binom{n+k}{n+1} = \frac{n+1-k}{n+1} \binom{n+k}{n}$. (For instance, see [6, p. 73].)

For their bijective proof, [2] used Krattenthaler's map: Ψ_K translates any right-to-left maximum of $\pi \in \mathcal{S}_n(132)$ into a return of the associated Dyck path. (A *return* of a Dyck path is a down-step landing on the x -axis.). The number of returns of Dyck paths is known to have a distribution given by $b(n-1, n-k)$. (This result was given in [4].)

Corollary 3.6 *The number of permutations $\pi \in \mathcal{S}_n(321)$ with $k-1$ elements $\pi_i = i+1$ is equal to $b(n-1, n-k)$.*

Proof. By the construction of $\Psi_{BJS}(\pi)$, a return, except the last (down-) step, appears if and only if i is an excedance of π with $\pi_i = i+1$. The very last step of $\Psi_{BJS}(\pi)$ is a return by definition. $\square$

Corollary 3.7 *The number of diagrams fitting in $(n-1, n-2, \dots, 1)$ with $k-1$ corners in the diagonal $i+j=n$ equals $b(n-1, n-k)$.*

Remark 3.8 In comparison with this, the Narayana number $N(n, k) = \frac{1}{n} \binom{n}{k-1} \binom{n}{k}$ counts the number of restricted diagrams with $k-1$ corners (see also Remark 2.6c)).

The condition $i+j=n$ on a corner (i, j) of the diagram $D(\pi)$ also occurs in the following section in context with the pattern 213 in a 132-avoiding permutation π .

We can also use the relation between diagrams and Dyck paths to obtain more information about the restricted partitions. For instance, the number of partitions in $\mathcal{Y}_n$ of a prescribed rank can easily be derived from a known result for paths. (The (*Durfee*) *rank* of a partition λ , denoted by $\text{rank}(\lambda)$, is the largest integer i for which $\lambda_i \geq i$, or equivalently, the length of the main diagonal of the diagram of λ .)

Lemma 3.9 *Let $\pi \in \mathcal{S}_n(132)$. With the above notation we have $w_{n+1}(\pi) = n - 2 \cdot \text{rank}(D(\pi))$.*

Proof. This is an immediate consequence of the relation between $D(\pi)$ and $\Psi_K(\pi)$. $\square$

Theorem 3.10 *Let $r(n, k)$ denote the number of partitions in $\mathcal{Y}_n$ of rank k . Then we have*

$$r(n, k) = \left(\frac{n+1-2k}{n+1-k} \binom{n}{k} \right)^2 \quad \text{for all } 0 \leq k \leq \lfloor \frac{n}{2} \rfloor.$$

Proof. Given a partition $\lambda \in \mathcal{Y}_n$ of rank k , let $\pi \in \mathcal{S}_n(132)$ be the (uniquely determined) permutation whose diagram equals the diagram of λ . By the lemma, both the first "half" and the second "half" of $\Psi_K(\pi)$ are paths from $(0, 0)$ to $(n, n-2k)$ only consisting of up-steps and down-steps and never falling below the x -axis (for the latter one, consider the reflection on the line $x=n$). Thus, $r(n, k)$ is the square of the number of precisely those paths. Feller's Ballot Theorem (see [6, p. 73]) enumerates the paths from the origin to (n, t) with $n, t \geq 1$ which neither touch (that is, there is no return) or cross the x -axis. Their number is equal to $\frac{t}{n} \binom{n}{\frac{n-t}{2}}$. Consequently, there are

$$\frac{t+1}{n+1} \binom{n+1}{\frac{n-t}{2}}$$

paths from $(0, 0)$ to (n, t) where $n, t \geq 0$ with up-steps and down-steps never going below the x -axis (returns are allowed). To see this, insert an up-step before the first step and define its starting point to be the origin. The resulting path satisfies the conditions of Feller's theorem. Conversely, each path from $(0, 0)$ to $(n+1, t+1)$ without returns can easily be transformed into an admissible path by deleting the first (up-) step and redefining the origin. Taking $t = n - 2k$ we obtain

$$r(n, k) = \left(\frac{n-2k+1}{n+1} \binom{n+1}{k} \right)^2 = \left(\frac{n+1-2k}{n+1-k} \binom{n}{k} \right)^2$$

for all $k = 0, \dots, \lfloor \frac{n}{2} \rfloor$. $\square$

Remark 3.11 By the proof, the positive root $q(n, k)$ of $r(n, k)$ counts the number of paths which begin at the origin, end at $(n, n - 2k)$, consist of up-steps and down-steps, and stay above the x -axis. Clearly, $q(n, 0) = 1$ and $q(n, \lfloor \frac{n}{2} \rfloor) = C_{\lceil \frac{n}{2} \rceil}$ for all n . Obviously, $q(n, k)$ satisfies the recurrence $q(n, k) = q(n - 1, k - 1) + q(n - 1, k)$. (The right-hand side is just the number of paths ending at $(n - 1, n - 2k + 1)$ and $(n - 1, n - 2k - 1)$, respectively.) Thus the numbers $q(n, k)$ are exactly the entries in the counter diagonals of the Catalan's triangle.

4 FORBIDDEN PATTERNS IN 132-AVOIDING PERMUTATIONS


Now we will use the correspondence between $\mathcal{S}_n(132)$ and $\mathcal{Y}_n$ for the enumeration of multiple restrictions on permutations. The results concerning the Wilf-equivalence of several pairs $\{132, \tau\}$ where $\tau \in \mathcal{S}_k$ are already known, see [12], [13], [9], [3]. (We say that $\{132, \tau_1\}$ and $\{132, \tau_2\}$ are *Wilf-equivalent* if $|\mathcal{S}_n(132, \tau_1)| = |\mathcal{S}_n(132, \tau_2)|$ for all n .) While the proofs given in these papers are analytical we present bijective ones.

Theorem 4.1 *Let $\pi \in \mathcal{S}_n(132)$ be a permutation not equal to the identity, and $k \geq 3$. Then*

- a) *π avoids $k(k - 1) \cdots 1$ if and only if $D(\pi)$ has at most $k - 2$ corners. In particular, we have $\text{des}(\pi) \leq k - 2$.*
- b) *π avoids $12 \cdots k$ if and only if $D(\pi)$ contains the diagram $(n + 1 - k, n - k, \dots, 1)$.*
- c) *π avoids $213 \cdots k$ if and only if every corner (i, j) of $D(\pi)$ satisfies $i + j \geq n + 3 - k$.*

Proof.

a)




Obviously, π contains a decreasing subsequence of length k if the diagram of π has at least $k - 1$ corners.

On the other hand, if there are at most $k - 2$ corners in $D(\pi)$, we have $\text{des}(\pi) \leq k - 2$ and hence $\pi \in \mathcal{S}_n(k \cdots 1)$. (Note that each corner of $D(\pi)$ corresponds to a descent in π .)

b) If the diagram $(n + 1 - k, n - k, \dots, 1)$ fits in $D(\pi)$ then we have $\pi_i \geq n + 3 - k - i$ for all i . Hence any increasing subsequence of π is of length at most $k - 1$: if $\pi_i = n + 3 - k - i$ then $i - 1$ elements from $i + k - 3$ many ones in $\{n + 4 - k - i, n + 5 - k - i, \dots, n\}$ appear in $\pi_1 \cdots \pi_{i-1}$. Conversely, let i be the smallest integer with $i + \pi_i < n + 3 - k$. Furthermore, choose j such that $\pi_j = n + 3 - k - i$ (by definition of i , we have $j > i$), and let $\pi_{i_1} < \dots < \pi_{i_{k-2}}$ be the elements of $\{n + 4 - k - i, n + 5 - k - i, \dots, n\}$ which are not equal to $\pi_1, \dots, \pi_{i-1}$. Note that

$j < i_1 < \dots < i_{k-2}$ since π is 132-avoiding. Thus $\pi_i \pi_j \pi_{i_1} \dots \pi_{i_{k-2}}$ is an increasing sequence.

c)



Let (i, j) be the top corner of $D(\pi)$ for which $i + j < n + 3 - k$. By removing the rows $1, \dots, i$ and the columns $\pi_1, \dots, \pi_i$, we obtain the diagram of a permutation $\sigma \in \mathcal{S}_{n-i}(132)$ whose letters are in the same relative order as $\pi_{i+1} \dots \pi_n$ where $\sigma_1 = \pi_{i+1} \leq j < \pi_i$. As discussed in the proof of part b), the element σ_1 is the first one of an increasing sequence of length $(n - i) - \sigma_1 + 1$ in σ . (Since $\sigma_1 \leq j < n + 3 - k - i$ the index $l = 1$ is the smallest one with $l + \sigma_l < (n - i) + 3 - k$.) Clearly, the first $j + 1 - \sigma_1$ terms are restricted by j . Thus there is an increasing sequence of length $n - (i + j) > k - 3$ in σ whose all elements are larger than π_i . Note that the elements $j + 1, j + 2, \dots, \pi_i - 1$ appear in $\pi_1 \dots \pi_{i-1}$. To prove the converse, suppose that every corner of $D(\pi)$ satisfies the condition given above. Then we have $\pi_i + i > n + 3 - k$ for all $i \in D(\pi)$. Hence for each descent i of π there exist at most $k - 3$ elements π_j with $j > i$ and $\pi_j > \pi_i$. Since π is 132-avoiding these elements form an increasing sequence. Thus there is no pattern $2134 \dots k$ in π . $\square$

Remarks 4.2

- a) From the statement of a), it follows that the maximum length of a decreasing subsequence of $\pi \in \mathcal{S}_n(132)$ is equal to the number of corners of $D(\pi)$ plus one, or in terms of permutation statistics, $\text{des}(\pi) + 1$. It is well-known that the length of the longest decreasing sequence can easily be obtained for any permutation in $\mathcal{S}_n$ via the Robinson-Schensted correspondence: it is just the number of rows of one of the tableaux P and Q corresponding to $\pi \in \mathcal{S}_n$.

It is clear from the construction that every left-to-right-minimum of π appears in the first column of P . Any other entry of this column must be the largest element of a 132-pattern in π . Hence in case of 132-avoiding permutations the elements of the first column of P are precisely the left-to-right-minima. As observed in Section 2, these minima are just the descent bottoms and the first letter of π . Thus for $\pi \in \mathcal{S}_n(132)$ the tableau P has $\text{des}(\pi) + 1$ rows.

- b) By part b), the length of a longest increasing subsequence of $\pi \in \mathcal{S}_n(132)$ equals the maximum value of $n + 1 - i - \lambda_i$ with $1 \leq i \leq n - 1$ where $\lambda_i \geq 0$ is the length of the i th row of $D(\pi)$. This also follows from Remark 2.6b) according to which λ_i is equal to the i th component of the code of π . Thus $n - i - \lambda_i$ counts the number of elements on the right of π_i which are larger than π_i . (Since π contains no pattern 132 these elements appear in increasing order.)

Corollary 4.3 $|\mathcal{S}_n(132, k(k-1) \dots 1)| = \frac{1}{n} \sum_{i=1}^{k-1} \binom{n}{i} \binom{n}{i-1}$ for all n and $k \geq 3$.

Proof. As mentioned in 2.6c) the number of partitions in $\mathcal{Y}_n$ whose diagram has exactly i corners is equal to the Narayana number $N(n, i+1) = \frac{1}{n} \binom{n}{i} \binom{n}{i+1}$. Thus there are $\sum_{i=0}^{k-2} N(n, i+1)$ diagrams with at most $k-2$ corners. $\square$

The following result also follows from a special case of [13, Th. 2.6].


Corollary 4.4 $|\mathcal{S}_n(132, 12 \cdots k)| = |\mathcal{S}_n(132, 213 \cdots k)|$ for all n and $k \geq 3$.

Proof. There is a simple bijection between the restricted diagrams which contain $(n+1-k, n-k, \dots, 1)$ and those ones whose all corners satisfy the condition $i+j \geq n+3-k$. (Note that the empty diagram associating with the identity in $\mathcal{S}_n$ belongs to the latter ones.) For each corner (i, j) of the diagram $(n+1-k, n-k, \dots, 1)$ we have $i+j = n+2-k$. Thus every diagram containing $(n+1-k, n-k, \dots, 1)$ is uniquely determined by its corners outside this shape (which are precisely the corners with $i+j \geq n+3-k$). Given such a diagram D , the corresponding diagram D' is defined to be this one whose corners are the corners of D which are not contained in $(n+1-k, n-k, \dots, 1)$. Conversely, for any diagram D' whose all corners satisfy $i+j \geq n+3-k$ we construct the corresponding diagram D as the union of D' and $(n+1-k, n-k, \dots, 1)$. $\square$

While the above conditions can be checked without effort, the characterization of the avoidance of the patterns considered now is more technical.

Given a permutation $\pi \in \mathcal{S}_n(132)$, let $\lambda_1, \dots, \lambda_l$ be the positive parts of the partition with the diagram $D(\pi)$. Let $a_i := n - (i + \lambda_i)$ for $i = 1, \dots, l$ and $b_i := n - (i + \lambda'_i)$ for $i = 1, \dots, \lambda_1$ where λ' denotes the conjugate of λ . Furthermore, for $i = 1, \dots, l$ let h_i be the length of the longest increasing sequence in $b_{\lambda_i} b_{\lambda_i-1} \cdots b_1$ whose first element is b_{λ_i} . We call the number h_i the *height of a_i* .

For example, the permutation $\pi = 8\ 9\ 5\ 4\ 6\ 7\ 2\ 3\ 10\ 1 \in \mathcal{S}_{10}(132)$ generates the diagram of $\lambda = (7, 7, 4, 3, 3, 3, 1, 1, 1)$:



The numbers $i + \lambda_i$ and $i + \lambda'_i$ are given on the left side and on the top of the diagram, respectively.

So we obtain $a(\pi) = (2, 1, 3, 3, 2, 1, 2, 1, 0)$ and $h(\pi) = (3, 3, 1, 2, 2, 2, 1, 1, 1)$. (Note that we have $b(\pi) = (0, 2, 1, 3, 3, 2, 1)$.)

Theorem 4.5 *Let $\pi \in \mathcal{S}_n(132)$ be a permutation, and let $a(\pi), h(\pi)$ be as above. Then π avoids the pattern $s(s+1) \cdots k12 \cdots (s-1)$, where $2 \leq s \leq k$ and $k \geq 3$, if and only if the longest decreasing subsequence of $a(\pi)$ which ends in an element of height $\geq s-1$ is of length at most $k-s$.*

Proof. Let i_1 be an integer such that $a_{i_1} \geq a_i$ for all $i < i_1$. Hence a decreasing sequence whose first element is a_{i_1} can not be extended to the left. Then $\lambda_{i_1} < \lambda_{i_1-1}$ where $\lambda_0 := n$. (Note that the condition $\lambda_i < \lambda_{i-1}$ is equivalent to $a_{i-1} \leq a_i$.) As shown in Section 2, the element π_{i_1} is a left-to-right minimum of π . Thus any increasing subsequence in π which starts with π_{i_1} is left maximal. In particular, we have $\pi_{i_1} = \lambda_{i_1} + 1$.

Now let a_j be an element of $a(\pi)$ with $a_j < a_{i_1}$, $j > i_1$, and $a_{i_1+1}, \dots, a_{j-1} > a_j$. Since $j-i > \lambda_i - \lambda_j$ for $i = i_1, i_1+1, \dots, j-1$, all the elements $\lambda_j+1, \lambda_j+2, \dots, \lambda_{i_1}$ occur in $\pi_{i_1+1}\pi_{i_1+2} \cdots \pi_{j-1}$. Hence $\pi_{i_1} < \pi_j$.

Consequently, if $a_{i_1} > a_{i_2} > \dots > a_{i_r}$ (with $i_1 < i_2 < \dots < i_r$) is a sequence such that there is no integer i with $i_l < i < i_{l+1}$ and $a_{i_l} > a_i \geq a_{i_{l+1}}$ for any l , then $\pi_{i_1} < \pi_{i_2} < \dots < \pi_{i_r}$ is an increasing subsequence of π which is maximal with respect to the property that π_{i_1} and π_{i_r} are its first and last elements, respectively. (Since π avoids the pattern 132, the relations $\pi_{i_1} < \pi_{i_l}$ for $l = 2, \dots, r$ imply that $\pi_{i_1}, \dots, \pi_{i_r}$ are increasing.)

It is clear from the definition that h_i is the maximal length of an increasing sequence of dots southwest of the dot representing π_i which begins at the top dot southwest of (i, π_i) . Thus, if a_{i_r} is an element of height $\geq s-1$ then there exist at least $s-1$ integers $i_r < j_1 < j_2 < \dots < j_{s-1}$ with $\pi_{j_1} < \dots < \pi_{j_{s-1}} < \pi_{i_r}$. Since π is 132-avoiding, we even have $\pi_{j_1} < \dots < \pi_{j_{s-1}} < \pi_{i_1}$. Choosing i_1 and i_r minimal and maximal, respectively, proves the assertion. $\square$

For any permutation $\pi \in \mathcal{S}_n(132)$ denote by $l_s(\pi)$ the largest integer l such that π contains the shifted pattern $s(s+1) \cdots l12 \cdots (s-1)$ where $s \geq 2$. By the theorem, $l_s(\pi)$ is equal to $s-1$ plus the maximum length of a decreasing sequence in $a(\pi)$ whose smallest element is of height at least $s-1$.

It is clear that the sequence $L(\pi) := (l_2(\pi) - 1, l_3(\pi) - 2, \dots)$ is a partition, that is, $l_s(\pi) + 1 \geq l_{s+1}(\pi)$ for all s . Since π avoids $s(s+1) \cdots k12 \cdots (s-1)$ if and only if no pattern $(k+2-s)(k+3-s) \cdots k12 \cdots (k+1-s)$ occurs in π^{-1} , the partition $L(\pi^{-1})$ is the conjugate of $L(\pi)$. (Obviously, for any permutation $\pi \in \mathcal{S}_n$ the diagram of the inverse π^{-1} is just the transpose of $D(\pi)$. Hence the set $\mathcal{S}_n(132)$ is closed under inversion.)

Remark 4.6 Using the relation between the diagram $D(\pi)$ and the Dyck path $\Psi_K(\pi)$, it is easy to see that the number a_i is just the height at which the i th down-step of $\Psi_K(\pi)$ ends. (We only consider the down-steps before the last up-step.) The numbers b_i needed for the construction of $h(\pi)$ are (in reverse order) the starting heights of the up-steps after the first down-step. Thus the

height h_i is precisely the number of (not necessarily consecutive) up-steps of increasing starting heights following the i th down-step in $\Psi_K(\pi)$. Hence, in case of $s = 2$ the theorem yields the second part of [9, Lemma Φ].

We shall prove now that the number of permutations in $\mathcal{S}_n$ which avoid both 132 and the pattern $s(s+1) \cdots k12 \cdots (s-1)$ with $k \geq 3$ and $s \in \{1, \dots, k\}$ does *not* depend on s .

Proposition 4.7 *Let $\pi \in \mathcal{S}_n(132)$, and let l be the maximum length of an increasing subsequence of π . Then π corresponds in a one-to-one fashion to a permutation $\sigma \in \mathcal{S}_n(132)$ with $l_2(\sigma) = l$.*

Proof. Let λ and μ be the partitions whose diagrams equal $D(\pi)$ and $D(\sigma)$, respectively. Given $\lambda = (\lambda_1, \dots, \lambda_{n-1}) \in \mathcal{Y}_n$, we define the sequence $\hat{\mu}$ by

$$\hat{\mu}_i := \begin{cases} \lambda_i + 1 & \text{if } \lambda_i + i < n \\ 0 & \text{if } \lambda_i + i = n \end{cases}$$

for $i = 1, \dots, n-1$, and obtain the partition μ by sorting $\hat{\mu}$. (Delete all parts $\hat{\mu}_i = 0$ with nonzero $\hat{\mu}_{i+1}$ and add the corresponding number of zeros at the end of the sequence.) It is obvious that $\mu \in \mathcal{Y}_n$, and it is easy to see that the map $\lambda \mapsto \mu$ is injective, and thus a bijection on $\mathcal{Y}_n$. To recover λ from μ , first set $\hat{\lambda}_i := \mu_i - 1$ for all positive μ_i . Then for any $\mu_i = 0$ let j be the largest integer for which $\hat{\lambda}_j + j \geq n-1$, and define $\hat{\lambda}$ to be the sequence obtained by inserting $n-1-j$ between $\hat{\lambda}_j$ and $\hat{\lambda}_{j+1}$. If there is no such an integer j then prepend $n-1$ to $\hat{\lambda}$. The partition resulting from this procedure is defined to be λ .

Consider now the sequence $\bar{a}(\pi) = (n-i-\lambda_i)_{i=1, \dots, n-1}$. (For the statement of Theorem 4.5 it suffices to consider the reduced sequence $a(\pi)$ which is obtained by omitting the final terms $\bar{a}_i = n-i$. By Remark 4.2b), we have $l = \max \bar{a}_i + 1$. Let j_1 be an integer satisfying $\bar{a}_{j_1} = l-1$. Using the definition of $\bar{a}(\pi)$ in terms of the Dyck path $\Psi_K(\pi)$, it is obvious that there exist some integers $j_1 < j_2 < \dots < j_{l-1} \leq n-1$ with $\bar{a}_{j_i} = l-i$ for $i = 1, \dots, l-1$. By Remark 4.6, the number $\bar{a}_i(\pi)$ is the height of the ending point of the i th down-step of $\Psi_K(\pi)$. (Note that the length l is equal to the maximum height of a peak of $\Psi_K(\pi)$.) By the construction, the elements of the sequence $a(\sigma)$ correspond to the nonzeros of $\bar{a}(\pi)$. In particular, we have $a_i(\sigma) > a_j(\sigma)$ for any $i < j$ if and only if the i th positive element of $\bar{a}(\pi)$ is larger than the j th positive one. Thus the elements corresponding to $\bar{a}_{j_1}, \bar{a}_{j_2}, \dots, \bar{a}_{j_{l-1}}$ form a decreasing sequence of maximal length in $a(\sigma)$. (For $l = 1$ we have $\lambda = (n-1, n-2, \dots, 1)$, i.e., $\pi = n(n-1) \cdots 1$ and hence $\mu = \emptyset$, i.e., $\sigma = \text{id} \in \mathcal{S}_n$.) Consequently, $l_2(\sigma) = (l-1) + 1$. (Note that any element $a_i(\sigma)$ is of height at least 1.) $\square$

The following result can be derived from the corresponding generating functions which were given for the first time by Chow and West ([3, Th. 3.1]). Several different analytical proofs appeared recently in [9, Th. 2, Th. 6] and [12, Th. 3.1].

Corollary 4.8 $|\mathcal{S}_n(132, 12 \cdots k)| = |\mathcal{S}_n(132, 23 \cdots k1)| = |\mathcal{S}_n(132, k12 \cdots (k-1))|$ for all n and $k \geq 3$.

Proof. The first identity is an immediate consequence of the preceding proposition. For the second one use that π avoids $23 \cdots k1$ if and only if π^{-1} contains no pattern $k12 \cdots (k-1)$. $\square$

Corollary 4.9 For any $s \geq 2$ there are as many partitions $\lambda \in \mathcal{Y}_n$ for which $i + \lambda_i \geq n + 2 - s$ for all i as such ones for which the sequence $a(\pi)$ for the corresponding permutation contains no element of height at least $s - 1$.

Proof. If $\lambda \in \mathcal{Y}_n$ satisfies the condition $i + \lambda_i \geq n + 2 - s$ for all i then its diagram contains $(n + 1 - s, n - s, \dots, 1)$. By Theorem 4.1b), this is equivalent to being $12 \cdots s$ -avoiding for the corresponding permutation. On the other hand, it follows from Theorem 4.5 (where $k = s$) that any permutation $\pi \in \mathcal{S}_n(132)$ avoids the pattern $s12 \cdots (s-1)$ if and only if every element of $a(\pi)$ is of height at most $s - 2$. Thus Corollary 4.8 proves the assertion. (Let τ be the partition associating with π . Then $\lambda \mapsto \mu \mapsto \tau := \mu'$ yields the desired bijection where $\lambda \mapsto \mu$ is the map appearing in the proof of 4.7, and μ' denotes the conjugate of μ .) $\square$

Now we can generalize the result from Proposition 4.7 for any $s \geq 2$.

Proposition 4.10 Let $\pi \in \mathcal{S}_n(132)$, and let $l \geq s - 1$ be the maximum length of an increasing subsequence of π . Then π corresponds in a one-to-one fashion to a permutation $\sigma \in \mathcal{S}_n(132)$ with $l_s(\sigma) = l$.

Proof. For $l \leq s - 1$ we have $i + \lambda_i \geq n + 2 - s$ for all parts λ_i of the partition corresponding to π . (Note that l is the maximum difference $n + 1 - (i + \lambda_i)$.) In this case, by Corollary 4.9, λ corresponds to a partition whose associated permutation satisfies $l_s = s - 1$. Thus we may assume that $l \geq s$.

The argument is completely parallel to that used to prove Proposition 4.7 but the construction of the bijection is more complicated. We preserve the notation of the proof of 4.7.

Given $\lambda = (\lambda_1, \dots, \lambda_{n-1}) \in \mathcal{Y}_n$, we define the sequence $\hat{\mu}$ by

$$\hat{\mu}_i := \begin{cases} \lambda_i + s - 1 & \text{if } \lambda_i + i < n + 2 - s \\ 0 & \text{if } \lambda_i + i = n + 2 - s \\ \vdots & \vdots \\ s - 2 & \text{if } \lambda_i + i = n \end{cases}$$

for $i = 1, \dots, n + 1 - s$, and $\hat{\mu}_i := \lambda_i$ otherwise. If there exists any integer i with $\hat{\mu}_i < s - 1$ and $\hat{\mu}_{i+1} \geq s - 1$ then sort $\hat{\mu}$ as follows: If $\hat{\mu}_i > 0$ then increase $\hat{\mu}_{i+1}$ by 1. Interchange $\hat{\mu}_i$ and $\hat{\mu}_{i+1}$. After this procedure we have $\hat{\mu} = \tau_1 \tau_2$ where τ_1 is a partition with parts at least $s - 1$,

and τ_2 is a subsequence containing elements of $\{0, \dots, s-2\}$ only. (Note that both τ_1 and τ_2 are nonempty for $s \geq 3$. Since $l \geq s$ there is an integer i satisfying $i + \lambda_i < n + 2 - s$ and hence $\hat{\mu}_i \geq s - 1$. On the other hand, $\hat{\mu}_{n+2-s}, \dots, \hat{\mu}_{n-1} \leq s - 2$ by definition.) It is clear from the construction that $i + \hat{\mu}_i \leq n$, and it is easy to see that the map $\lambda \mapsto \hat{\mu}$ is injective. To recover λ from $\hat{\mu}$, first define a sequence $\hat{\lambda}$ by $\hat{\lambda}_i := \hat{\mu}_i + 1 - s$ for all indices i with $\hat{\mu}_i \geq s - 1$ and $\hat{\lambda}_i := \hat{\mu}_i - i + n + 2 - s$ otherwise where $i = 1, \dots, n + 1 - s$. If there exists any integer with $\hat{\lambda}_i < \hat{\lambda}_{i+1}$ then decrease $\hat{\mu}_i$ by 1 if $\hat{\mu}_{i+1}$ is positive, interchange $\hat{\mu}_i$ and $\hat{\mu}_{i+1}$, and determine $\hat{\lambda}$ again for the resulting sequence $\hat{\mu}$. (This procedure will be done while $\hat{\lambda}$ is not a partition.) Adding $\hat{\mu}_{n+2-s}, \dots, \hat{\mu}_{n-1}$ at $\hat{\lambda}$ completes the partition λ .

If τ_2 , and hence $\hat{\mu}$, is a partition then set $\mu := \hat{\mu}$. The partitions $\lambda \in \mathcal{Y}_n$ for which this case occurs are characterized by the following conditions:

- (i) $\lambda_{n+2-s} = \dots = \lambda_{n-1} = 0$.
- (ii) If $i + \lambda_i \geq n + 2 - s$ for any $i \leq n - s$ then $\lambda_{i+1} < \lambda_i$.

Suppose that there exists an integer $i \geq n + 2 - s$ with $\hat{\mu}_i = \lambda_i > 0$. By Remark 4.6, the i th down-step of $\Psi_K(\pi)$ ends at a point of height $n - (i + \lambda_i) \leq n - (n + 2 - s + \lambda_i) = s - 2 - \lambda_i \leq s - 3$. Since $l \geq s$ there is an integer j with $j + \lambda_j < n + 2 - s$. (Obviously, $j < n + 2 - s \leq i$.) The ending point of the corresponding down-step is of height at least $s - 1$. Thus there must exist an integer k with $j < k < i$ such that $k + \lambda_k = n + 2 - s$, that is, $\hat{\mu}_k = 0$. The second condition is evident. (In case $s = 2$ the above conditions are satisfied for all partitions in $\mathcal{Y}_n$.)

If τ_2 is not ordered then we obtain the partition μ by increasing all elements of $\hat{\mu}$ which are smaller than $s - 1$ and smaller than an element to their right.

Consider the sequences $\bar{a}(\pi)$ and $a(\sigma)$ again. By the construction, each element of $\bar{a}(\pi)$ which is at least $s - 1$ corresponds to a term a_i of $a(\sigma)$ whose height is at least $s - 1$ or for which there exists an element a_j with $i < j$ and $h_j(\sigma) \geq s - 1$. (Then necessarily $a_i \geq a_j$.) In particular, we have $a_{i_1}(\sigma) > a_{i_2}(\sigma)$ for any $i_1 < i_2 \leq i_3$ where $h_{i_3}(\sigma) \geq s - 1$ if and only if the corresponding elements in $\bar{a}(\pi)$ are in the same order. Analogously to the proof of Proposition 4.7 this yields the assertion. $\square$

An analytical proof of the following result was given in [13, Th. 2.4].

Corollary 4.11 $|\mathcal{S}_n(132, 12 \cdots k)| = |\mathcal{S}_n(132, s(s+1) \cdots k12 \cdots (s-1))|$ for all n and $k \geq 3$ and $2 \leq s \leq k$.

Proof. This follows immediately from the previous proposition. $\square$

5 A FINAL NOTE

In Section 2 we have shown that the diagram of a permutation indicates the existence of some

subsequences of type 132. But in positive case, we even obtain the exact number of occurrences.


In [8], Fulton defined the following rank function on the essential set. Given a corner (i, j) of the diagram $D(\pi)$, i.e. $(i, j) \in \mathcal{E}(\pi)$, its *rank* is defined by the number of dots northwest of (i, j) and denoted by $\rho((i, j))$.

It is clear from the construction that the number of dots in the northwest is the same for all diagram squares which are connected. (This yields the fundamental property of the ranked essential set of a permutation π , that it uniquely determines π .) So, we can extend the rank function on $D(\pi)$. The information about the number of sequences of type 132 containing in a permutation is encoded by the ranks of its diagram squares.

Theorem 5.1 *Let $\pi \in \mathcal{S}_n$ be a permutation, and let $D(\pi)$ be its diagram. Then the number of occurrences of the pattern 132 in π is equal to*

$$\sum_{(i,j) \in D(\pi)} \rho((i, j)).$$


Proof. Extending the arguments of Theorem 2.2, it is easy to see that each square (i, j) of $D(\pi)$ corresponds to exactly $\rho((i, j))$ subsequences of type 132 in π , namely the sequences $k \pi_i j$ where k ranges over all column indices of dots northwest of (i, j) :



□

Remark 5.2 As mentioned above, we have $|D(\pi)| = \text{inv}(\pi)$ for all $\pi \in \mathcal{S}_n$. Hence the non-weighted sum $\sum_{(i,j) \in D(\pi)} 1$ counts the number of occurrences of the pattern 21 in π .

Example 5.3 The ranked diagram of $\pi = 4\ 2\ 8\ 3\ 6\ 9\ 7\ 5\ 1\ 10 \in \mathcal{S}_{10}$ is



Thus π contains 20 subsequences of type 132 and 18 inversions.

In [14], Mansour and Vainshtein studied the generating function for the number of permutations on n letters containing exactly $r \geq 0$ occurrences of pattern 132.

Acknowledgement I am grateful to Toufik Mansour for pointing me to some literature relating to restricted permutations.

REFERENCES

1. S. C. Billey, W. Jockusch and R. P. Stanley, Some Combinatorial Properties of Schubert Polynomials, *J. Alg. Comb.* **2** (1993), 345-374.
2. P. Brändén, A. Claesson and E. Steingrímsson, Catalan continued fractions and increasing subsequences in permutations, *Discrete Math.* **258** (2002), 275-287.
3. T. Chow and J. West, Forbidden subsequences and Chebyshev polynomials, *Discrete Math.* **204** (1999), 119-128.
4. E. Deutsch, A bijection on Dyck paths and its consequences, *Discrete Math.* **179** (1998), 253-256.
5. K. Eriksson and S. Linusson, Combinatorics of Fulton's essential set, *Duke Math. J.* **85** (1996), 61-80.
6. W. Feller, An Introduction to Probability Theory and Its Applications, vol. I, Wiley, New York, 1968.
7. M. Fulmek, Enumeration of permutations containing a prescribed number of occurrences of a pattern of length 3, preprint math.CO/0112092, 2001.
8. W. Fulton, Flags, Schubert polynomials, degeneracy loci, and determinantal formulas, *Duke Math. J.* **65** (1992), 381-420.
9. C. Krattenthaler, Permutations with restricted patterns and Dyck paths, *Adv. Appl. Math.* **27** (2001), 510-530.
10. I. G. Macdonald, Notes on Schubert Polynomials, LaCIM, Université du Québec à Montréal, 1991.
11. T. Mansour and A. Vainshtein, Restricted permutations and Chebyshev polynomials, *Sémin. Lothar. Comb.* **47** (2002), B47c.
12. T. Mansour and A. Vainshtein, Restricted permutations, continued fractions, and Chebyshev polynomials, *Electron. J. Comb.* **7** (2000), R17.
13. T. Mansour and A. Vainshtein, Restricted 132-avoiding permutations, *Adv. Appl. Math.* **26** (2001), 258-269.
14. T. Mansour and A. Vainshtein, Counting occurrences of 132 in a permutation, *Adv. Appl. Math.* **28** (2002), 185-195.
15. A. Reifegerste, Differenzen in Permutationen: Über den Zusammenhang von Permutationen, Polyominos und Motzkin-Pfaden, Ph.D. Thesis, University of Magdeburg, 2002.

16. A. Reifegerste, The excedances and descents of bi-increasing permutations, in preparation.
17. R. Simion and F. W. Schmidt, Restricted Permutations, *Europ. J. Combinatorics* **6** (1985), 383-406.
18. J. West, Generating trees and the Catalan and Schröder numbers, *Discrete Math.* **146** (1995), 247-262.